

ISSN 0869-2904

EXPERIMENT

IN Mineralogy Crystallography
Petrology Geophysics
Geochemistry Isotopy

GeoSciences

2020 Volume 26 Number 1



EXPERIMENT

in

Mineralogy
Petrology
Geochemistry

Crystallography
Geophysics
Isotopy

GeoSciences

Founded in 1991 by

INSTITUTE of
EXPERIMENTAL
MINERALOGY

*Russian Academy of
Sciences*

EXPERIMENT IN GEOSCIENCES is the only journal in English that gives you the latest, most reliable information on advances in experimental mineralogy, petrology, geochemistry, crystallography and geophysics in Russia. The coverage of the journal includes:

- original research and review articles
- abstracts of scientific papers
- proceedings, abstracts and announcements of national and international conferences, symposia and workshop
- information about the leading experimental centers in Russia
- advertisement of new techniques and equipment

Editor in Chief
O.G. Safonov

Editors:
T.F.Kazachenko
tanya@iem.ac.ru

Ass.Editors@Comp.graphics:
M.S.Belova
belova@iem.ac.ru

Editorial Board

V. Yu. Chevychelov
D.A.Chareev
L.Lakshtanov
O.A.Lukanin
E.G.Osadchii
Yu.N.Pal`yanov
A.F.Redkin
Yu.B.Shapovalov
N.I.Suk
M.V.Voronin

Institute of Experimental Mineralogy

Russian Academy of Sciences
142432 Chernogolovka
Academica Osip'yana str, 4,
Moscow region
Russian Federation

<http://www.iem.ac.ru>

TRANSLATED into ENGLISH by:
the authors

Copyright (C) IEM

Photographic reproduction, microfilm, electronic data base, video disks, or any other reproduction of text, figures, or tables from this journal is prohibited without the publishers' permission

CONTENTS

Problems of Planetology, Cosmochemistry and Meteoritics 5

Alexeev V.A., Pavlova T.A., Kalinina G.K. Some features of the radiation history of ureilites.....	5
Badekha ¹ K. A., Grokhovsky ² V.I. Phase transformation in the mono crystals of meteoritic and laboratory origin. UDC 669.017.3	8
Barenbaum A.A. ¹ , Shpekin M.I. ² Cumulative hypothesis of formation on the moon of the south pole-aitken basin by galactic comets UDC: 550.2: 523.4-1/-8	11
Barenbaum A.A. ¹ , Titorenko A. ² Optimized galactic model of geological cyclicity and its testing on geology and astronomy data UDC 550.93, 551.7, 523.525, 524.66	15
Fisenko A.V., Semjonova L.F. Alternative composition of xenon components in nanodiamonds of meteorites. UDC 550.42	19
Ivanov A.A., Sevastyanov V.S., Shnykin B.A., Dolgonosov A.A., Krivenko A.P., Priymak S.V., Galimov E.M. Evaluation of the various energy sources effectiveness at the stage of prebiological structures formation during abiogenesis. UDC 550.47	23
Kuyunko N.S., Alexeev V.A. Thermoluminescent research and determination of metamorphism degree of carbonaceous CO and CV group chondrites.....	25
Lavrentjeva Z.A., Lyul A.Yu. To question about formation of the pallasite parent bodies: on the peculiarities of trace element composition of mineral components from the omolon pallasite.....	28
Lyul A.Yu., Lavrentjeva Z.A. On an influence of the metamorphic processes on siderophile element distributions between the major components of the h-group of ordinary chondrites.....	31
Nikitin S.M. ¹ , Gorbatshevich F.F. ² , Skripnik A.Ya. ³ , Beltyukov N.A. ⁴ , Morozov I.A. ⁴ Brittle fracture features of terrestrial analogues of ordinary chondrites components under volumetric compression	33
Shornikov S. I. ¹ , Yakovlev O. I. ¹ The chemical fractionation of perovskite at evaporation	36
Tselmovich V.A. ¹ , Maxe L.P. ² Microstructure and composition of particles of native iron of cosmic origin UDC 523.68140	43
Yurkovets V.P. General morphological features of some fragments of the Ladoga space body UDC 551.4:552.6	43
Zharkova E.V., Lukanin O.A., Tsekhonina T.I., Senin V.G. Intrinsic oxygen fugacity measurements of impact glasses from impact craters El'gygytgyn (Chukotka, Russia) and Zhamanshin (Kazakhstan) UDC 550.843 (543.559)	45

Mineral equilibria at high PT-parameters 50

Bulatov V.K. ^{1,3} , Girmis A.V. ^{2,3} , Brey G.P. ³ , Woodland A. ³ , Höfer H. ³ Ferropericlasite crystallization under upper mantle conditions.....	50
Fedkin V.V., Sipavina L.V. Clinopyroxene in the eclogite-blueschist complexes: structural signs of the conditions of its formation UDC 549.6+552.16:552.48.....	54
Gorbachev N.S., Kostyuk A.V., Nekrasov A.N., Gorbachev P. N., Sultanov D.M. The interaction of phlogopite with carbonate at $P = 4$ GPa, $T = 1200-1300$ °C: phase relationships and stability of phlogopite. UDC 550.4.02	58
Gorbachev N.S., Kostyuk A.V., Gorbachev P.N., Sultanov D.M. Distribution of elements between phlogopite and carbonate at $P = 2.8$ GPa, $T = 1250$ °C. UDC 550.4.02.....	61
Iskrina A.V. ^{1,2} , Spivak A.V. ² , Bobrov A.V. ^{1,2,3} , Dubrovinsky L.S. ⁴ , Eremin N.N. ¹ , Marchenko E.I. ¹ . Experimental study of calcium-ferrite structure type phases in the mg-al-cr-o system at conditions of the earth's transition zone and lower mantle UDC 549.02	63
Kogarko L.N. Crystallization of alkaline phosphates from magmatic melt. UDC 553.491.8:549.271(571.5).....	65
Kostyuk A.V., Gorbachev N.S., Sultanov D.M., Nekrasov A.N. Experimental investigation of peridotite-basalt-(K, Na) ₂ CO ₃ -H ₂ O system at $P = 4$ GPa, $T = 1400$ °C: phase composition and critical relationship between melt and fluid. UDC 550.4.02	68

Thermodynamic properties of minerals and fluids 72

Korepanov Ya.I., Osadchii E.G. Thermodynamic properties of Ag-Au alloy with high concentration of silver ($0.9 < x < 1$ Ag _x Au _{1-x}).....	72
Shornikov S. I. ¹ , Ivanova M.A. ¹ , Minaeva M. S. ² Thermodynamic properties of the MgO – FeO melts	74
Shornikov S. I. Thermodynamic properties of the CaO–MgO melts	77

Synthesis of minerals 81

- Gurbanova O.A.¹, Evdokimov A.I.¹, Antipin A.M.², Volkov A.S.¹, Khasanova S.³, Dimitrova O.V.¹ Hydrothermal synthesis and crystal structure of a new orthophosphate of lithium, aluminum, copper and iron. UDC 548.736..... 81
- Kovalskaya T.N.¹, Ermolaeva V.N.¹, Varlamov D.A.¹, Kalinin G.M.¹, Kovalskiy G.A.^{1,2}, Chaychuk K.D.¹ Synthesis of eudialyte from alkaline pegmatites. Preliminary data. 82

Hydrothermal equilibria and ore formation 85

- Alekseyev V.A., Burmistrov A.A., Gromiak I.N. Quartz dissolution in water vapor and solution distillation in quartz–water–vapor system at 300°C..... 85
- Balitsky V.S., Balitskaya L.V., Golunova M.A., Setkova T.V., Bublikova T.M. New experimental data on phase transformations and states of synthetic water-hydrocarbon inclusions in quartz 88
- Bublikova T.M., Setkova T.V., Balitsky V.S. Theoretical study of the solubility of basic copper carbonates in ammonium chloride systems 91
- Korzhinskaya V.S., Kotova N.P. Temperature effect on the behavior of Ta and Nb during the dissolution OF pyrochlore, tantalite, β -Ta₂O₅ and β -Nb₂O₅ in (HF+HCl) solutions..... 93
- Kotelnikov A.R.¹, Daminov B.B.², Daminova L.B.², Bryanskiy N.V.³, Akhmedzhanova G.M.¹, Suk N.I.¹ Synthetic fluid inclusions in quartz: a check for the adequacy of capture of ore elements. UDC 550.4.02 95
- Kotelnikov A.R.¹, Suk N.I.¹, Kotelnikova Z.A.^{1,2}, Korzhinskaya V.S. Fluids in equilibrium with silicate substance and problems of ore genesis UDC 550.89:553.062..... 97
- Redkin A.F. Tungsten valence in hydrothermal solutions on the experimental and calculated data. 98
- Sidkina E.S.¹, Mironenko M.V.¹, Polyakov V.B.² Calculation of serpentinization of olivin. Verification of the equilibrium-kinetic model. 100

The formation and differentiation of magmas 105

- Bortnikov N.S.¹, Kryazhev S.G.², Gorelikova N.V.¹, Smirnov S.Z.³, Gonevchuk V.G.⁴, Semenyak B.I.⁴, Dubinina E.O.¹, Sokolova E.N.³ The fluid regime of Badzhal hydrothermal-magmatic system of the Far East (Priamurie, Russia)..... 105
- Persikov E.S., Bukhtiyarov P.G., Shekleina M.D. Features of basalt melt crystallization at moderate hydrogen pressures (preliminary results). 111
- Rusak A.A.¹, Shehekina T.I.², Alferyeva Ya.O.², Gramenitskiy Eu.N.², Zinovieva N.G.², Khvostikov V.A.³, Kotelnikov A.R.⁴ Features of phase crystallization in a high-fluorine model granite system at a temperature drop from 700 TO 400 °C and a pressure of 1 kbar UDC 552.11, 550.42. 114
- Tobelko D.P.¹, Portnyagin M.V.^{1,2}, Krashennikov S.P.¹ Complex methodology for estimating the primary H₂O content in island-arc magmas (the case of Kamchatka)..... 117

Physical chemical properties of geomaterials 120

- Setkova T.V.¹, Spivak A.V.¹, Zakharchenko E.S.¹, Borovikova A.Yu.², Nesterova V.A.^{1,2}, Balitsky V.S.¹, Bublikova T.M.¹, Pushcharovsky D.Yu.² Raman spectroscopy of synthetic Ga,Ge-rich tourmaline at high pressure up to 30 Gpa (preliminary data)..... 120
- Zharikov A.V.¹, Lebedev E.B.², Rodkin M.V.³ Anomalies of rock physical properties due to phase transitions and their possible role in seismotectonic processes..... 121

Experimental geocology 126

- Grishantseva E.S.^{1*}, Alekhin Yu.V.¹, Drozdova O.Yu.¹, Demin V.V.², Zavgorodnyaya Y.A.² Experimental studies of organic matter in natural waters of lakes in Vladimir region using a set of analytical methods. UDC 550.4.02 126
- Salavatova D.S., Fiaizullina R.V., Bychkov D.A. Adsorption extraction of mercury ions from an aqueous solution by rocks of the Maykop series. UDC 550.424: 550.4.02: 550.41: 550.42 129

Engineering of experiment 133

CONTENTS

Chertkova N.V. ¹ , Spivak A.V. ¹ , Zakharchenko E.S. ^{1,2} , Litvin Y.A. ¹ , Kuzyura A.V. ¹ , Safonov O.G. ¹ , Efimchenko V.S. ² , Meletov K.P. ^{1,2} . A refinement of the externally heated diamond anvil cell technique for laboratory applications, UDC 53.086; 608.2; 543.424.2	133
Lakshantov L.Z., Karaseva O.N. Study of the mechanisms of chalk recrystallization. UDC 550.4.02	134
Molchanov ¹ V.P., Medkov ² M.A. Ways of industrial development of gold-ilmenite placers of the sikhote-alin with the use of pyro- hydrometallurgical methods	137

Interaction in the systems of fluid–melt–crystal 140

Alferyeva Ya.O., Novikova A.S., Gramenitskiy E.N. Interaction of fluorine-containing granite melt and calcite as a possible cause of formation of high-calcium ongonites. UDC 552.113	140
Bychkov D.A., Koptev-Dvornikov E.V., Romanova E.S. Verification of the system of thermobarometers-compositometers on experimental and natural material. UDC 552.111: 550.41	141
Chevychelov V.Yu. On the solubility of ferrotapiolite and nb/ta ratios in model high-aluminous, subaluminous and alkaline granitoid melts UDC 550.42	144
Chevychelov V.Yu., Viryus A.A. Solubility of ilmenorutile in model granitoid melts at various alkalinity–alumina. UDC 550.42	147
Khodorevskaya L.I. Experimental modelling of alkaline metasomatism under pressure gradient conditions at 750°C. UDC 553.065.1	149
Koptev-Dvornikov E.V., Romanova E.S., Bychkov D.A. Orthopyroxene liquidus thermobarometer-compositometer for a range of melt compositions from magnesian basites to dacites	152
Kotelnikov A.R., Korzhinskaya V.S., Suk N.I., Van K.V. Experimental study of zircon and hafnon solubility in silicate melts UDC 550.89	155
Kotelnikov A.R. ¹ , Suk N.I. ¹ Damdinov B.B. ² , Damdinova L.B. ² , Khubanov V.B. ² Experimental studies of phenakite solubility in silicate melts UDC 550.89	158
Novikova A.S., Alferyeva Ya. O., Gramenitskiy E.N. Phase relationship modelling in syrostan massive near-contact zone by 1 kbar and 900 and 850 °C. UDC 552.113	159
Romanova E.S., Koptev-Dvornikov E.V., Bychkov D.A. Pigeonite liquidus thermobarometer for a range of melts compositions from magnesian basites to dacites. UDC 552.111: 550.41	161
Seliutina N.E. ^{1,2} , Safonov O.G. ^{2,1} , Varlamov D.A. ² Petrological and experimental study of syenitization of tonalite gneisses exemplified by the midiapala massif, limpopo complex, South Africa. UDC 552.11	164
Shchekina T.I. ¹ , Zinovieva N.G. ¹ , Rusak A.A. ² , Khvostikov V.A. ³ , Gramenitskiy E.N. ¹ , Alferyeva Ya.O. ¹ , Kotelnikov A.R. ⁴ Particularities of scandium distribution between silicate and salt melts and crystal phases in the Si-Al-Na-K-Li-F-O-H system AT 800°-500°C and 1 kbar. UDC 552.11, 550.42	166
Suk N.I., Kotelnikov A.R., Viryus A.A. Experimental studies of loparite solubility in silicate melts	169

AUTHORS INDEX 172

Experimental geocology

Grishantseva E.S.^{1,*}, Alekhin Yu.V.¹,
Drozdova O.Yu.¹, Demin V.V.²,
Zavgorodnyaya Y.A.² **Experimental studies
of organic matter in natural waters of lakes in
Vladimir region using a set of analytical
methods.** UDC 550.4.02

¹ Department of Geology, M.V. Lomonosov Moscow State University, Moscow

² Department of Soil Science, M.V. Lomonosov Moscow State University, Moscow

* Correspondence: shes99@mail.ru

Abstract. Organic matter in natural waters takes an active part in the migration of trace elements, playing a crucial role in the biogeochemical cycles of trace elements. The paper discusses the results of research on the organic matter of natural waters of lakes in Vladimir region, obtained using a set of analytical methods. Selected as objects of the study were representative waterbodies with different levels of primary productivity: eutrophic lake Ershevik, dystrophic lake Ignatkovo. The proposed set of analytical methods for determining the content, composition and forms of finding organic matter and related trace elements includes determining the content of dissolved organic carbon and N on the liquid element analyzer LiquiTOC trace, Elementar; studying the molecular mass distribution of organic matter by high-performance liquid chromatography in a variant of gel-penetrating chromatography; studying the forms of finding organic matter and trace elements by cascade filtration with the separation of size fractions; spectrophotometric measurement of water density using a Specord 50 (ANALYTIK JENA) and 511 UV/Vis (Portlab) spectrophotometer to study the absorption spectra of samples in the UV and visible regions of the spectrum, followed by the calculation of indicators that characterize the dissolved organic matter, the degree of its humification, the proportion of autochthonous and allochthonous organic matter.

Keywords: *organic matter; natural waters; complex of analytical methods.*

The aim of the research was to study, typify and carry out a comparative analysis of the organic matter in natural waters of the lakes of Vladimir Oblast. Identification of regional and local differences in the migration of ionic forms, organic complexes, and suspensions in natural waters is of great practical importance for prognostication and justification of quantitative models of river runoff evolution under the influence of various natural and anthropogenic factors. To obtain information on sustainable correlations between microelements and organic matter, as well as the mechanisms of their concerted transport and deposition in various natural conditions, as objects of study we chose reservoirs with different levels of primary productivity: the eutrophic Lake Ershevik and the dystrophic Lake Ignatkovo in Vladimir Oblast in the drainage basin of the Klyazma River. The water of Lake Ershevik is characterized by near neutral pH values of about 7,

while in Lake Ignatkovo pH is equal to 5.7. Lake Ershevik had higher indices of chromaticity, permanganate oxidizability and BOD₅: chromaticity of -179 degrees of Cr-Co scale; permanganate oxidizability of 20 mg O₂/dm³; biological oxygen demand of 2 mg O₂/dm³.

Organic matter in natural waters takes an active part in the processes of migration of trace elements, playing a decisive role in the biogeochemical cycles of trace elements. The work discusses the results of studying the organic matter in natural waters of the lakes of Vladimir Oblast obtained using a complex of analytical methods. As objects of study, representative water bodies with different levels of primary productivity were selected: the eutrophic Lake Ershevik and the dystrophic Lake Ignatkovo. Separately, we analyzed the waters of the surface euphotic horizon, the bottom horizon, and silt waters extracted from bottom sediments.

To study the distribution of organic matter and trace elements in natural waters by size fractions and to separate truly dissolved forms of metals from suspended and colloidal particles in natural waters, we applied the method of cascade membrane filtration and ultrafiltration with pore diameters of 0.45 μm, 0.22 μm and 0.014 μm (1 kDa) using an Amicon device, followed by an analysis of the content of a large number of trace elements in the isolated fractions of natural waters by the ICP-MS method (Element 2, Thermo Scientific) (Alekhin, 2010). In samples of natural waters, hydrochemical indices were determined according to standard GOST methods. The proposed complex of analytical methods for determining the content, composition and forms of organic matter and associated trace elements also includes the determination of the content of dissolved organic carbon (DOC) and N on the liquid elemental analyzer LiquiTOC trace (Elementar), spectrophotometric measurement of water density on a Specord 50 spectrophotometer (ANALYTIK JENA) and 511 UV/Vis spectrophotometer (Portlab) to analyze the absorption spectra of samples in the UV and visible regions, and study of the molecular weight distribution of organic matter by high-performance liquid chromatography (specifically, gel permeation chromatography) on an Agilent 1100 chromatograph with a diode array detector. Column: TSK G2000SW 7.5x300 mm. Eluent: 0.1 M phosphate buffer, pH 7 + 0.1% SDS (sodium dodecyl sulfate). Speed: 0.5 ml/min. Sample volume: 50 μl. Based on the results obtained, we calculated indices that characterize the dissolved organic matter, degree of its humification, proportions of autochthonous and allochthonous organic matter.




Fig.1. Absorption spectra of dissolved organic matter in natural waters.



Fig.2. Diagrams of the mass distribution of organic matter and trace elements by size fractions of natural waters (for example, oz.Ershevik).




Fig.3. Chromatogram of molecular mass distribution of organic matter of natural waters (filtrate 0.22 μm).

The absorption spectra (Fig. 1) were used to calculate the indices characterizing the dissolved organic matter: SUVA, E_{254}/E_{436} . The SUVA index was calculated as the ratio of the absorption at 254 nm to the content of dissolved organic carbon in the sample (Table 1). The value of $SUVA > 4$ indicates a predominance of hydrophobic and especially aromatic compounds, while $SUVA < 3$ corresponds to predominantly hydrophilic compounds (Edzwald, Tobiason 1999). The E_{254}/E_{436} ratio is calculated in order to assess the predominance of autochthonous or allochthonous dissolved organic carbon (Battin, 1998). Lake Ershevik had higher values of optical density that correlate with the degree of humification of dissolved organic matter; for Lake Ignatkovo, this index was 2 times lower. The predominant role of the

hydrophobic and aromatic fraction in the composition of organic matter is characteristic of Lake Ershevik, especially in the summer, whereas the dystrophic Lake Ignatkovo is characterized by the predominance of the hydrophilic fraction of organic matter. The high values of the E_{254}/E_{436} ratio in Lake Ershevik, especially in the bottom horizon and silt waters, confirm the predominant role of autochthonous organic matter formed as a result of the biogeochemical activity of aquatic plants and plankton inside the reservoir. For the dystrophic Lake Ignatkovo, the indices were found to be lower, which indicates a significant contribution of terrigenous organic matter in the composition of dissolved organic matter.

Table 1. Main characteristics of organic matter of natural waters of lakes of the Vladimir region.

	DOC, mg/l	N, mg/l	E_{254}	E_{436}	C/N	SUVA	E_{254}/E_{436}
Lake Ershevik							
unfiltered	19.3	1.09	0.868	0.053	18	5	16
surface water (< 0.22 μm)	18.3	0.98	0.698	0.048	19	4	15
surface water (< 1 kDa)	16.5	0.89	0.496	0.021	19	3	24
bottom waters (< 0.22 μm)	16.4	0.97	0.540	0.029	17	3	19
bottom waters (< 1 kDa)	14.7	0.99	0.392	0.013	15	3	30
silt water (< 0.22 μm)	6.6	4.70	0.123	0.005	1.4	2	25
silt water (< 1 kDa)	5.7	4.56	0.095	0.002	1.3	2	48
Lake Ignatkovo							
unfiltered	9.9	0.51	0.306	0.019	19.4	3	16
surface water (< 0.22 μm)	9.3	0.39	0.295	0.021	23.8	3	14
surface water (< 1 kDa)	8.8	0.32	0.243	0.011	27.5	3	22
bottom waters (< 0.22 μm)	10.0	1.03	0.36	0.029	9.7	4	12
bottom waters (< 1 kDa)	7.0	0.62	0.179	0.008	11.3	3	22
silt water (< 0.22 μm)	8.7	1.59	0.187	0.018	5.5	2	10
silt water (< 1 kDa)	6.3	0.97	0.114	0.010	6.5	2	11

The study of the distribution of organic matter in natural waters by size fractions indicates the predominance of low molecular weight organic matter (<1kDa), the proportion of which reaches 98% (Fig. 2). Not more than 39% of dissolved organic carbon is represented by a fraction from 0.22 μm to 0.014 μm (1 kDa). The distribution of organic matter and trace elements by fractions is identical for the lakes under study and allows to identify a general pattern of the behavior of trace elements that is characteristic of ecosystems of water bodies in the Central Russia. A comparative analysis of the organic matter in surface waters showed that small lakes are characterized by rather high concentrations of dissolved organic carbon (9-19 mg/L).

The results of the study of the molecular weight distribution of organic substances by HPLC showed that the chromatograms obtained for surface waters

are characterized by a monomodal distribution and have an identical form. In the surface waters of both lakes, the composition of organic matter is predominated by low molecular weight organic compounds with average molecular weights $M_w=269$ Da for Lake Ignatkovo and $M_w=258$ Da for Lake Ershevik (Fig. 3). Moreover, their concentration in Lake Ignatkovo is significantly higher than that in Lake Ershevik.

The statistical treatment of analytical data by the method of factorial and correlation analysis using the Gold-Geochemist program allowed to identify several groups of elements according to the degree of correlation of their content with the content of dissolved organic carbon, Fe and Mn.




Fig.4. Dendrogram of correlations.

The proposed complex of analytical methods makes it possible to carry out a comprehensive study of the content, composition and forms of organic matter and associated trace elements to identify the patterns of their concerted migration in aquatic ecosystems.

The work was supported by the Russian Foundation for Basic Research (Grants No. 19-05-00519, 18-05-00162).

References

- Alekhin Yu. V., Ilina S. M., Lapitsky S. A., Sitnikova M. V. Results of studying the joint migration of trace elements and organic matter in the river flow of the boreal zone// Vestn. Mosk. UN-TA. Ser. GEOL. 2010. no. 6, p. 49-55. In-Rus.
- Battin T.J. Dissolved organic materials and its optical properties in a blackwater tributary of the upper Orinoco River, Venezuela. *Organic Geochemistry*, 28, 1998, - p.561-569.
- Edzwald J.K., Tobiason J.E. Enhanced coagulation: US requirements and a broader view. *Water Science and Technology*, 40, 1999. – p.63-70.

Salavatova D.S., Fiaizullina R.V., Bychkov D.A. Adsorption extraction of mercury ions from an aqueous solution by rocks of the Maykop series. UDC 550.424: 550.4.02: 550.41: 550.42

Lomonosov Moscow State University, Geological Faculty, Department of Geochemistry, Russia, Moscow (salavatova-jamilya2012@yandex.ru, fiaizullina@geol.msu.ru, krok@geol.msu.ru)

Abstract. The behavior of mercury in the mud volcanic process has not been sufficiently studied. Mercury accumulates on clays by adsorption from water, and then desorption occurs and it begins its own behavior in the

mud volcanic system. The determination of sorption characteristics for clays of the Maykop series of varying degrees of transformation has become our first priority. In this work, we studied the dependence of mercury adsorption on the pH of the solution and the duration of the contact of the solution with the adsorbent. Experimental adsorption isotherms are obtained, which are analyzed using the equations of Langmuir, Freundlich, Redlich-Peterson, and Toth. It has been shown that the observed adsorption process is described by the Redlich-Peterson and Toth models, which indicates the predominantly chemical nature of adsorption and that the adsorption centers are not identical in their properties.

Keywords *mercury adsorption; mercury in mud volcanoes; adsorption isotherms; Maykop series.*

Mud volcanism is provided by two processes: the process of transformation of sedimentary rocks or the so-called illite-smectite transition, i.e. the conversion of mixed-layer clays with a high water content to hydromica minerals such as illite with a lower water content, the process of water evolution and the sorption properties of clay minerals change: their specific surface decreases and, accordingly, the components that were in the sorbed complex of mixed-layer smectite minerals transfer to the newly formed water. The behavior of mercury under these conditions has not been adequately studied. Mercury accumulates on clays by adsorption from water, and then its desorption occurs and it begins its own behavior in the mud volcanic system. Little is said about the behavior of mercury in the processes of mud volcanism in modern literature. For example, in the work of Portuguese colleagues (Mieiro, Pato et al., 2007), cores of sediments of a mud volcano in the Gulf of Cadiz (Atlantic Ocean, Portugal, Spain) were studied and it was shown that the mercury content in them varied from 9.0 to 86.2 ppb. There are generalized data on the mercury content in the waters of mud volcanoes of the Kerch Peninsula, the North-

West Caucasus, Azerbaijan and Turkmenistan, which ranged from 0.2 to 0.5 ppm, with an average concentration of 0.38 ± 0.19 ppm, calculated on the basis of 10 measurements (Ivanov, 1996). Thus, the aim of this study was to study the sorption properties of clays of the Maykop series of varying degrees of conversion with respect to mercury (II) ions. The laboratory air temperature during all experiments was $25.6 \pm 2.3^\circ\text{C}$. To measure the equilibrium concentration of mercury, the “cold vapor” method was used with an atomic absorption spectrometry using Portable Mercury Analyzer PMA-1 (EcON, Moscow) with a PAR-3m attachment. The reducing agent was a 1% sodium borohydride solution in a 1% sodium alkali solution. To control the acidity of the solutions, the Expert-001 pH meter (Econix-Expert, Moscow) was used. As the pH electrode, the EGK-10601 combined glass electrode (Measuring equipment, Moscow) was used. The concentration of adsorbed mercury was determined by the difference in concentrations in solutions according to the equation (Fiaizullina et al., 2020):

$$q_e = \frac{(C - C_e) \cdot m_{sol}}{m_{sorb}}, \quad (1)$$

where q_e is the amount of Hg(II) adsorbed at equilibrium is (ppb); C and C_e are the initial and equilibrium concentrations of metal ions in solution (ppb); m_{sol} is the mass of the solution (g); m_{sorb} is the mass of adsorbent (g). The degree of mercury

extraction (E , %) from solutions was calculated by the equation:

$$E = \frac{C - C_e}{C} \cdot 100\%. \quad (2)$$

The effect of pH. It was shown that the maximum adsorption of mercury occurs at $\text{pH} = 3-4$ (Salavatova, Fiaizullina, 2019) and is 240 ppm ($E = 99.3\%$). When discussing changes in the degree of extraction of elements by clays in an acidic environment, the dissolution of clays cannot be ruled out. However, in our opinion, the dissolution of clay components in a medium with $\text{pH} = 3-4$ is unlikely, since the clay samples we study mainly contain clay minerals and quartz, which are highly acid resistant and should not dissolve under these conditions. All experiments described in this work were performed in the range of 3-4 pH units.

Kinetics of adsorption. To study the kinetics of adsorption on clay of the Maykop series of low degree of conversion, two series were delivered: with and without shaking. In a series with stirring, the tubes were shaken manually 3 times a day. The exposure time ranged from 15 minutes to 1 month. Figure 1 shows that after 4 days under dynamic conditions the mercury removal completeness reaches more than 96% - the process reaches equilibrium. Under conditions without shaking, equilibrium occurs only after 2 weeks and persists after a month.




Fig. 1. The degree of mercury extraction over time (there is the square root of time in hours on the abscissa)

Adsorption isotherms. Analysis of adsorption isotherms gives an idea of the features of the adsorption process, the effectiveness of the selected sorbent and allows us to evaluate the feasibility of its use for the absorption of certain substances. In the present work, adsorption isotherms were analyzed

using the equations of Langmuir (Langmuir, 1916), Freundlich (Freundlich, 1906), Redlich-Peterson (Redlich, Peterson, 1959), and Toth (Toth, 1971). Traditionally, to find the parameters of equations, their linearized forms are used. However, as shown in (Boulinguez et al., 2008), it is much more

Experimental geocology

efficient to minimize the nonlinear chi-square test with solver add-in function of the Microsoft Excel. All further results were obtained using this method. The values of the parameters of the equations and the χ^2 are given in table 1. The chi-square test was calculated according to the equation:

$$\chi^2 = \sum_{i=1}^n \frac{(q_e^{calc.} - q_e^{meas.})^2}{q_e^{meas.}}, \quad (3)$$

where χ^2 is nonlinear chi-square test, q_e^{calc} is calculated adsorbate concentration at equilibrium (mg/g), q_e^{meas} is measured adsorbate concentration at equilibrium (mg/g). Small χ^2 value indicates a good description using the selected adsorption model.

Table 1. The adsorption isotherm parameters calculated using four models and χ^2

Isoterm	Equation	Parameter	Value	χ^2
Langmuir	$q_e = q^\infty \cdot \frac{K_L \cdot C_e}{1 + K_L \cdot C_e}$	q^∞ , mg/g	9.01	2.08
		K_L	0.00147	
Freundlich	$q_e = K_F \cdot C_e^{\frac{1}{n}}$	n	2.08	0.677
		K_F	0.118	
Redlich-Peterson	$q_e = \frac{K_R \cdot C_e}{1 + a_R \cdot C_e^g}$	g	0.641	0.241
		a_R	0.113	
		K_R	0.0415	
Toth	$q_e = \frac{K_T \cdot C_e}{(a_T + C_e)^t}$	t	1.62	0.208
		a_T	38.3	
		K_T	0.284	




Fig. 2. Isotherms of mercury adsorption on clay of the Maykop series calculated according to the Langmuir and Freundlich equations

Figure 2 shows adsorption isotherms calculated according to the Langmuir and Freundlich equations and experimental points. It can be seen from the obtained data that the Langmuir isotherm poorly describes the adsorption process. This is due to the fact that the Langmuir equation assumes single-layer

homogeneous adsorption with a limited number of equivalent adsorption centers, where each center has a constant enthalpy and sorption activation energy. The Freundlich isotherm describes the adsorption process well in the region of high concentrations, but is completely inapplicable in the region of low

concentrations, for which it has been criticized several times.

The most accurately studied adsorption process is described by the Redlich-Peterson and Toth models (fig. 3), as evidenced by the lowest values of χ^2 . This indicates the predominantly chemical nature of adsorption and that the adsorption centers are not identical in their properties. The Redlich-Peterson isotherm (fig. 3a) was developed as a hybrid model of the Langmuir and Freundlich isotherms. The Toth isotherm (fig. 3b) was developed to improve the approximation of the experimental data by the Langmuir isotherm. It is useful for describing

heterogeneous adsorption systems and satisfactorily describes equilibria at both low and high concentrations of the adsorbent. The form of the model equation assumes an asymmetric quasi-Gaussian energy distribution, with most adsorption centers having an adsorption energy below the average value. The results obtained show that clays of the Maykop series of a low degree of conversion actively adsorb mercury from aqueous solutions. The maximum adsorption is more than 10 mg/g (1 wt.%), which suggests that a significant amount of mercury can accumulate in these clays.




Fig. 3. Isotherms of mercury adsorption on clay of the Maykop series calculated according to the equations of Redlich-Peterson (a) and Toth (b)

Thus, it was found that when the mass ratio of the sorbent and the solution is 1: 500, the adsorption equilibrium occurs after 4 days with active shaking and after 2 weeks without shaking. At this point, the completeness of removal reaches 96% and 94%, respectively. With a longer exposure (1 month), equilibrium is maintained. The experimental data on mercury adsorption are best described by the Redlich-Peterson and Toth equations.

- Langmuir I. The constitution and fundamental properties of solids and liquids // *J. Am. Chem. Soc.*, 1916. Vol. 38(11). P. 2221-2295.
- Miêro C.L., Pato P., Pereira E., Mirante F., Coutinho J.A.P., Pinheiro L.M., Magalhães V.H., Duarte A.C. Total mercury in sediments from mud volcanoes in Gulf of Cadiz // *Marine Pollution Bulletin*, 2007. Vol. 54. P. 1539-1544.
- Redlich O., Peterson D.L. A useful adsorption isotherm // *J. Phys. Chem.*, 1959. Vol. 63. P. 1024-1026.
- Toth J. State equations of the solid gas interface layer // *Acta Chem. Acad. Hung.*, 1971. Vol. 69. P. 311-317.

This work was supported by RSF grant №19-17-00200 and RFBR grant №19-05-00519.

References

- Ivanov V.V. Ecological geochemistry of elements. Ref. book, in 6 books. Book 5. M.: Nedra, 1996. – 352 p.
- Salavatova D.S., Fiaizullina R.V. Modeling of Mercury (II) Adsorption on Clays of the Maykop Series // *Proceedings of the X All-Russian School of Young Scientists Experimental Mineralogy, Petrology and Geochemistry. IEM RAS Chernogolovka*, 2019. – P. 122-124.
- Fiaizullina R.V., Kuznetsov E.V., Salavatova D.S. Adsorption of Mercury from an Aqueous Solution on a Synthetic Organosilicon Sorbent // *Mos. Univ. Geol. Bull.*, 2020. Series 4: Geology. In Print.
- Boulinguez B., Le Cloirec P., Wolbert D. Revisiting the determination of Langmuir parameters application to tetrahydrothiophene adsorption onto activated carbon // *Langmuir*, 2008. Vol. 24. P. 6420-6424.
- Freundlich H.M.F. Over the adsorption in solution // *J. Phys. Chem.*, 1906. Vol. 57. P. 385-470.