

European Psychiatry

THE JOURNAL OF THE EUROPEAN
PSYCHIATRIC ASSOCIATION

**EPA 2021
ABSTRACT
SUPPLEMENT**



EUROPEAN
PSYCHIATRIC
ASSOCIATION



CAMBRIDGE
UNIVERSITY PRESS



CONTENTS

Abstracted in: Science Citation Index (SCI) - Clarivate, Social Science Citation Index (SSCI) - Clarivate, Scopus, Medline/
PubMed, Google Scholar, PsycINFO

Abstracts of the 29th European Congress of Psychiatry - 2021

Plenary	S1
State of the Art	S2
Debate	S4
Joint Symposium	S6
Symposium	S8
Workshop	S47
ECP Programme	S65
Oral Communication	S74
e-Poster Presentation	S211
e-Poster Viewing	S763

hospitalization in 25,6%. Fetal health problems were detected in 7,2%. A psychiatric trouble has been reported by 4% of the pregnancies. The total score of PAI ranged from 27 to 82 in our sample. We found a statistically significant negative correlation between PAI and duration of marriage ($p=0,012$); PAI and gestation number ($p=0,039$); and a correlation between PAI and the planning of the pregnancy ($p=0,030$).

Conclusions: Socio-demographic and clinical factors should be taken in consideration while evaluating pregnant women at risk of perinatal psychological difficulties.

Conflict of interest: No significant relationships.

EPP0087

Prevalence of adhd among adolescents with type 1 diabetes mellitus: Meta-analysis of observational studies.

J. Bhela^{1*}, M. Amin², R. Subhedar³, J. Gude⁴ and B. Sheikh⁵

¹Psychiatry, Case Western Reserve / Metrohealth hospital, Cleveland, United States of America; ²Psychiatry, Windsor University School of Medicine, Frankfurt, United States of America; ³Psychiatry, McMaster University, Brampton, Canada; ⁴Psychiatry, Northwell Health/Loong Island Jewish Hospital, Queens, United States of America and ⁵Psychiatry, Brookdale Hospital Medical Center, Brooklyn, United States of America

*Corresponding author.

doi: forthcoming

Introduction: Prevalence of type 1 diabetes mellitus (T1DM) and attention deficit hyperactivity disorder (ADHD) in adolescents has been increasing remarkably every year and the impact of T1DM on ADHD prevalence has been rarely explored.

Objectives: The aim of the study was to evaluate the prevalence of ADHD in adolescents with T1DM.

Methods: Advanced literature search on PubMed was carried out following MOOSE guidelines to identify the observational studies conducted on comparing the prevalence of ADHD in adolescents with or without T1DM using MeSH-terms x OR y from December 1, 2019, to April 30, 2020. Seven studies were identified out of 2,300 search results with the consensus of 3 reviewers. Five studies were selected for data analysis where three studies reported positive outcomes and two reported no association between ADHD and T1DM. Comprehensive meta-analysis software with mixed effects random models was used to analyze pooled OR and 95%CI and created forest plot.

Results: Out of five studies with data on the prevalence of ADHD in T1DM, three studies had details on positive outcomes. Pooled prevalence of ADHD was higher amongst adolescents with T1DM [1.85% vs 1.33%; 95%CI; $p<0.001$; 386/20,900 vs 31,732/2,384,121 adolescents; Heterogeneity (I²): <0.0001]. In compare to non-T1DM, T1DM patients' ADHD had higher odds [aOR:1.62; 95%CI:1.45-1.79; $p<0.00001$; I²:0.000; $p=0.42$].

Conclusions: Close attention, proper management and treatment is required for adolescents with Type 1 Diabetes Mellitus in order to reduce the higher prevalence of ADHD in this age group. The association between T1DM between ADHD has rarely been studied and needs to be investigated further.

Keywords: ADHD and diabetes; prevalence of ADHD in type 1 diabetes; ADHD and type 1 diabetes in adolescents; ADHD and type 1 diabetes

EPP0088

Conditions of negation formation in children of early age with down syndrome

E. Sedova^{1*}, V. Stepanova² and Z. Gardanova¹

¹Psychological-social Department, Pirogov Russian National Research Medical University, Moscow, Russian Federation and ²Psychology, Charitable Fund Downside Up, Moscow, Russian Federation

*Corresponding author.

doi: forthcoming

Introduction: According to Vygotsky, children with special needs follow the same trajectory of development as normally developing children, although some of the skills can be observed in a later period. This statement can be implemented to the children with Down syndrome. The number of such children in Russia is around 25 thousand. **Objectives:** The aim is to study the conditions of negation formation in children with Down syndrome.

Methods: The sample consisted of 22 dyads of children with Down syndrome of 24-36 months old and their mothers. The research methods included: parents' questionnaire; analysis of problematic situations; Tkacheva's inventory Parent's Psychological Type; Varga & Stolin Inventory of Parental Attitude; Toronto Alexithymia Scale, Bass-Darky Hostility Questionnaire, Leonhard-Schmieschek Test, Spielberger's Test Anxiety Inventory.

Results: Firstly, we have studied how a child expresses his or her negative reaction: whether he or she uses a gesture or a sound for "no" or reacts with the whole body. According to those results we have divided the sample into two groups and then have compared them. The research shows the connection between mother's aggressiveness and formation of the child's negation reaction (gesture/sound or the whole body) as well as differences in the level of alexithymia and anxiety: all the characteristics are lower in the first group.

Conclusions: Mothers of the children with Down syndrome demonstrate a high and a medium level of anxiety. However, the mothers of the children who expresses negation with a gesture/sound show a lower anxiety level comparing with the mothers whose children react with the whole body.

Keywords: Down syndrome; parent-child relations; negation; children with special needs

EPP0089

Productivity of tasks performance in children and adolescents with different level of media multitasking

A. Koshevaya*, G. Soldatova and S. Chigarkova

Faculty Of Psychology, Lomonosov Moscow State University, Moscow, Russian Federation

*Corresponding author.

doi: forthcoming

Introduction: Media multitasking (MMT) begins to play an increasingly important role in terms of the digitalization of everyday life. At the same time, the effect of MMT on efficiency is still poorly highlighted, especially among the younger generation (May, Elder, 2018; Patterson, 2017; Peifer, Zipp, 2019).

Objectives: The aim is to identify types of MMT in children and compare them by productivity and time of task performance.

Methods: Quasi-experimental research, which included the performance of tasks on a computer and a smartphone, was conducted among children in three groups aged 7-10 years, 11-13 years, 14-16 years (N=154).

Results: Based on a number of criteria, several types of behavior are identified in the MMT environment: two subgroups of “single-taskers” (23%), “light” (19%), “medium” (54%) and “heavy” (4%) MMT. Comparative analysis ($p=0.027$) reveals high scores for proper task performance of the subgroup “single-taskers1”, as well as “heavy” and “light” MMT. The other subgroup “single-taskers2” and “medium” MMT show, on the contrary, low productivity results. MMT groups also differ in terms of task performance time ($p=0.006$). The “light” and “heavy” MMT cope the fastest. The second place by the speed of task performance is held by “medium” MMTs and “single-taskers1”, and the slowest are “single-taskers2”.

Conclusions: The results show that both linear and nonlinear behavioral strategies can lead to a certain level of performance. Most children try to operate in the MMT mode and it is becoming the dominant and ubiquitous modus vivendi for the younger generation. The reported study was funded by RFBR, project No. 19-29-14181.

Conflict of interest: The reported study was funded by RFBR, project No. 19-29-14181.

Keywords: media multitasking; productivity; Children; adolescents

EPP0090

Early premorbid of children with ASD

M. Kalinina^{1*} and G. Shimonova²

¹Child Psychiatry, MHRC RAMS, Moscow, Russian Federation and

²Child Psychiatry, MHRC RAMS, Moscow, Russian Federation

*Corresponding author.

doi: forthcoming

Introduction: According to numerous studies, the origins of most endogenous mental diseases, in particular, autism, lie in early childhood. This concept is based on the classical theories of diathesis.

Objectives: The purpose of this study was to clarify the phenomenology of mental disorders in children with an assessment of the prognostic significance of symptoms that preceded the development of the disease for future mental health.

Methods: The clinical material was collected during the survey of a child population. Selected for prospective observation was 40 children (1-3 years old) from the high-risk group for schizophrenia with functional disorders of the endogenous spectrum. All patients were examined by clinical methods and pathopsychological, neuropsychological. Psychometric scales PANSS, CARS were used. The results were mathematically evaluated using the Statistica 7 program.

Results: The clinical picture of the mental state of young children, in children with autistic disorders in 1,5-3 years, was determined by a specific complex of disorders, which were reduced to a general deficit, especially in the emotional sphere, vegetative dysregulation, most often, the sleep-Wake rhythm. Motor skills, as a rule, did not lag significantly behind the age standards. The onset of actual autistic disorders was noted older than 1 year of life. Children received medication and corrective therapy. The detailed clinical picture of violations developed gradually. Dynamics of psychopathological picture in (80,0%) children was regressive.

Conclusions: The study shows the importance of preventive measures in people related to ASD, sparing individual approach in education and therapy.

Keywords: early childhood; autism; high risk group; a prospective study; autism; early childhood

EPP0091

Observing flaming or trolling online: Prevalence in Russian youth and adolescents and relationship to tolerance and aggression

G. Soldatova, S. Chigarkova, A. Koshevaya and E. Rasskazova*

Faculty Of Psychology, Lomonosov Moscow State University, Moscow, Russian Federation

*Corresponding author.

doi: forthcoming

Introduction: Cyberaggression including the most wide-spread variants of flaming (O’Sullivan, Flanagan, 2003; Voggeser et al., 2017) and trolling (Buckels et al., 2018) is affecting mental health of adolescents and youth although it could be (Kowalski, 2014; Wright, Wachs, 2020).

Objectives: The aim was to study prevalence of flaming and trolling experience in Russian youth and adolescents and its relationship to general aggression and tolerance.

Methods: 525 adolescents 12-13 years old, 1029 adolescents 14-17 years old, 736 youth 18-30 years from 8 Federal regions in Russia appraised their experience of flaming or trolling online (as initiators, victims and observers) using vignettes. 1105 parents of adolescents appraised whether their children experienced flaming or trolling online. Then they filled Aggression Questionnaire (Buss, Perry, 1992) and Tolerance Index (Psychodiagnostics..., 2008).

Results: More than one-half of adolescents (51-58% in 12-13 years old and 64% in 14-17 years old) and youth (45-69%) reported experience of flaming and trolling online, mostly as observers (32-65%). Parents accurately appraised flaming experience in their children but underestimated trolling experience ($p<.05$). Adolescents and youth observing flaming online report higher hostility, anger and physical aggression ($F=17.8-28.3$, $p<.01$, $\eta^2=.02$) while lower social tolerance ($F=4.27$, $p<.05$, $\eta^2=.01$). In adolescents observing trolling online these effects are stronger than in youth observing trolling online (interaction: $F=5.68$, $p<.05$, $\eta^2=.01$).

Conclusions: Observing trolling and flaming online is related to higher aggression and low tolerance in adolescents and youth and for adolescents the relationship is stronger. The reported study was funded by RFBR, project 20-013-00857.

Conflict of interest: The reported study was funded by RFBR, project 20-013-00857.

Keywords: adolescents; cyberaggression; flaming; trolling

EPP0092

Neurocognitive functions in children and adolescents with different enthusiasm for video games

G. Soldatova¹, A. Vishneva^{2*} and A. Koshevaya¹

¹Faculty Of Psychology, Lomonosov Moscow State University, Moscow, Russian Federation and ²Clinical Psychology, center for speech pathology and neurorehabilitation, Moscow, Russian Federation

*Corresponding author.

doi: forthcoming

Introduction: Video games are becoming increasingly popular among children (Lenhart et al., 2015). There is a lack of research that studies the impact of online games on children’s neurocognitive functions.